

Human Resources

South Cove Community Health Center

Infectious Medical Waste Program

Presents:



Infectious Medical Waste Information

Guidance on Segregation of Wastes
and Waste Reduction

Why Infectious Waste is Regulated

- To minimize the potential for the spread of disease from a medical setting to the general public; and
- To reduce the overall amount of infectious medical waste produced.
 - helps to protect the environment, and reduces medical facilities' treatment expenditures

Diseases Caused by Bloodborne Pathogens

- HIV / AIDS
- Hepatitis B

- Arboviral infections
- Brucellosis
- Creutzfeldt-Jakob Disease
- Hepatitis C
- Leptospirosis
- Malaria
- Rabies
- Syphilis
- Tularemia
- Viral Hemorrhagic Fevers

What is Infectious Medical Waste

Infectious Medical Waste is defined as medical waste capable of producing an infectious disease.

Waste is considered Infectious when it is:

- Contaminated by an organism that is pathogenic to healthy humans;
- The organism is not routinely available in the environment; and
- The organism is in significant quantity and virulence to transmit disease.

Infectious Wastes Specifically Are

- Blood and blood products in a free flowing, unabsorbed state;
- Contaminated sharps,
- Isolation Wastes,
- Laboratory wastes, and
- Unfixed pathological tissues





Black Bag
Black Bag on floor
CNCI



Infectious Laboratory Wastes



- Cultures
- Etiological agents
- Specimens
- Stocks
- Related contaminated wastes
- Vaccine vials



Pathological Wastes

- Fixed Pathological wastes are not Infectious Medical Waste
- Unfixed Pathological wastes must be incinerated
 - wastes containing pathological items must be appropriately labeled to ensure they are incinerated



Infectious Isolation Wastes

- Wastes generated from the care of a patient who has or is suspected of having a disease caused by a CDC Class 4 agent, listed below

CLASS 4 VIRAL AGENTS:

- Alastrim, Smallpox, Monkey pox, and White pox.
- Hemorrhagic fever agents, including Crimean hemorrhagic fever (Congo), Junin, and Machupo viruses
- Herpes virus simiae (Monkey B virus)
- Lassa virus
- Marburg virus
- Tick-borne encephalitis virus complex, including Russian spring-summer encephalitis, Kyasanur forest disease, Omsk hemorrhagic fever, and Central European encephalitis viruses
- Venezuelan equine encephalitis virus
- Yellow fever virus

Disposal of Isolation Wastes

- Isolation wastes that do not meet the definition of infectious medical waste should be separated and disposed in the general waste stream
 - disposable gowns
 - face masks
 - shoe covers

All waste from an isolation room should be treated with caution and the appropriate Personal Protective Equipment (PPE) must be worn during handling and disposal.

What about OSHA

- Ensure that all employees can safely perform their normal duties without undue health risks
- Bloodborne Pathogen (BBP) Standard developed to protect employees with occupational exposure to bloodborne pathogens
 - HIV
 - HbV
- Employers required to evaluate engineering controls to reduce or eliminate employee exposure risks
 - adoption of a needleless system

Bloodborne Pathogen Standard

Employers must:

- Ensure that Universal Precautions are observed
- Provide free Hepatitis-B vaccination series
- Provide all necessary PPE and ensure that it is used
- Provide BBP training at hire, and annually thereafter
- Maintain records of all training
- Maintain an Exposure Control Plan, update annually
- Record exposure incidents and follow-up activities

Bloodborne Pathogen Standard

Defines Infectious Medical Waste as:

- Liquid or semi-liquid blood or other potentially infectious materials (OPIM),
- Contaminated items that would release blood or OPIM in a liquid or semi-liquid state if compressed,
- Items caked with dried blood or OPIM that would dislodge during handling,
- Contaminated sharps, and
- Pathological and microbial wastes containing blood or OPIM

Other Potentially Infectious Material

OPIIM

- Any body fluid with visible blood
- Amniotic fluid
- Cerebrospinal fluid
- Pericardial fluid
- Peritoneal fluid
- Pleural fluid
- Saliva in dental procedures
- Semen/vaginal secretions
- Synovial fluid
- Anywhere body fluids are indistinguishable



Infectious Waste Is Not

- Used personal hygiene products
 - tissues
 - feminine products
 - diapers
- Gauze and dressings containing small amounts of blood,
- Fixed pathological tissues,
- Uncontaminated medical tubing and devices

Tubing with any visible fluid blood must be disposed in the biohazard waste



Preventing Disease Transmission

- The single most effective measure to control the transmission of Bloodborne Pathogens is:

Universal Precautions

- Treat all human blood and other potentially infectious materials like they are infectious for Hepatitis B and HIV

Blood and Fluid Borne Pathogen Exposures

Exposures to blood and fluid borne pathogens in the medical setting typically occur by one of the following ways:

- Puncture from contaminated needles, broken glass, or other sharps
- Contact between non-intact (cut, abraded, acne, or sunburned) skin and infectious body fluids
- Direct contact between mucous membranes and infectious body fluids

Example: A splash in the eyes, nose, or mouth

Transmission of Blood and Fluid Borne Pathogens

An exposure incident does not guarantee disease transmission. Several factors affect transmission:

- Infected Source – the disease stage of the source
- Means of Entry - severity or depth of the: puncture wound, broken skin, or direct contact with mucus membrane
- Infective Dose - the amount and type of fluid, as well as the amount of infectious agent in the fluid
Blood is the fluid of greatest concern
- Susceptible Host - immunocompromised

Prevention of Exposure

Guidelines to reduce the risk of exposure:

- ★ Frequent hand washing
- Use of standard barrier precautions
- Regular cleaning and decontamination of work surfaces with a cleaning agent labeled as effective against Mycobacterium/TB
- Vaccination against Hepatitis-B
- Proper infectious waste disposal

Exposure Incident Response

- Wash exposed area with soap and water
- Flush splashes to nose, mouth or skin with water
- Irrigate eyes with water or saline
- Report the exposure to supervisor
- Follow your facility's exposure response plan
- Report all exposures, regardless of severity, for your safety

Collection of Infectious Waste

- Infectious medical wastes must be collected at the point of generation in the appropriate color coded bags
- Biohazard bags must be labeled with the international biohazard symbol and appropriate wording; “biohazard,” “biomedical waste,” “infectious medical waste,” or “regulated medical waste”



Sharps

- Must be collected at the point of generation, in a leak-proof and puncture-resistant container
- Containers must bear the international biohazard symbol and appropriate wording



- Containers should never be completely filled, nor filled above the full line indicated on box.

Liquid Infectious Medical Wastes

Liquid Infectious Medical Waste, i.e., the contents of suction canisters, may be disposed of in several ways:



- Placed directly in the Biohazardous waste,
- Poured down a sanitary sewer,
- Solidified using an approved disinfectant solidifier and discarded in the solid waste

Packaging and Storage

Wastes shall be collected in a lined, cardboard box or reusable plastic container that is labeled with the biohazard symbol and appropriate wording.

- Once the box or container is full, the bag lining it must be sealed and the container then sealed shut
- Boxes must be labeled with facility name, address, phone and fax numbers, and the date
- A full, sealed container can be stored on site for no more than 30 days

Shipping and Manifests

Every load of waste shipped off-site for destruction is tracked using a manifest system

- The manifest is a multiple copy document that accompanies the waste to the treatment facility
- Every individual who takes possession of the waste, including someone from your facility, must sign the manifest
- As the waste generator, your facility is responsible for the waste until you receive the proof-of-destruction copy of the manifest

Tracking Waste Through Destruction

- The proof-of-destruction copy of the manifest must be returned to your facility within 50 days from the date that the load of waste was picked up.
- Massachusetts Infectious Waste Program must be notified of any load that has not been accounted for within 50 days.
- All proof-of-destruction manifests must be kept on file at your facility for 3 years.

This manifest has been signed by the treatment facility, certifying that this load has been destroyed and is no longer infectious.

Manifest form with various sections including:

- Header information (date, time, location)
- Signature of the transporter
- Signature of the treatment facility representative
- Facility certification section with a red arrow pointing to the signature.
- Company information: **RAMSONG, INC.**, 3212 RIDGE ROAD, MATTHEWS, NC 28105, 704-841-4746

Bloodborne Pathogen Spill Kit

All medical facilities must have a spill kit. All employees should know where the spill kit is located.

It must contain the following items:

- Absorbent material to manage ten gallons of fluid
- One gallon hospital grade disinfectant and a spray bottle
- Many large biohazard bags
- 2 sets of disposable coveralls, boots, caps
- 2 pair of heavy neoprene gloves
- 2 pair of eye and respiratory protection devices
- 100 yards of boundary marking tape, 1 roll of packing tape
- First aid kit (facilities with an emergency room exempt)

Over Classification

- The improper disposal of solid wastes that do not meet the definition of infectious medical waste, as if they were infectious
- It is the most commonly cited violation, with 98% of permitted facilities being marked
- It increases the financial burden on patients and taxpayers in the form of increased disposal costs for health care facilities







Problems with Over Classification

- When large volumes of plastics, common in medical waste, are incinerated there is an increased potential for atmospheric release of carcinogenic agents
- Increased medical waste generation increases the risk of costly accidents and spills due to the increased number of trucks required to haul the waste



Why Bother Segregate Wastes?

Costs for waste disposal:

- About \$0.01 per pound to haul regular waste to a landfill.
- From \$0.28, and increases to over \$4.00 per pound to haul and treat as infectious waste.



A Real Life Example

During an inspection of one southern West Virginia facility it was noted that the ER, ICU, Lab, OB, and OR departments were not segregating any waste.

All waste was being disposed of as infectious.

Management said the staff was too busy to segregate garbage.

- In 1996 this facility generated 245,060 lbs. of infectious waste.
- They paid \$0.29 cents per pound to have their infectious waste hauled away and treated.

The Cost of Over Classification

- The inspection revealed 90 - 95% over classification.
- The total cost for disposal of infectious waste for 1996 was \$71,067.40
- Cost savings potential of \$64,000 to \$68,000




Teamwork Made it Possible

This same facility two years later:

- In 1998 they generated 114,000 pounds.
- A reduction of about 50%.
- In 2001 they generated 58,838 pounds.
- A total reduction of more than 76%
- This facility realized a \$54,004 cost savings

Source Separation is the Key!

- Everyone needs to consider which waste stream an item goes in every time wastes are disposed
- We realize that you are busy, but it only takes a few seconds to separate waste items into the proper waste stream
- By properly segregating medical wastes, the weight of infectious waste can be drastically reduced in every facility





Routinely Over Classified Items

- Diapers (adult and baby)
- Paper towels
- Unsaturated dressings and chucks
- Wrappers and packaging
- IV bags and oxygen tubing
- Gloves with no visible contamination
- Urine catheters and bags
- Paper, newspapers, and food containers
- Urine cups and specimen containers with no visible blood
- Empty Medication vials and broken glass











INFECTIONIOUS WASTE
Do not reuse sharps in
recycling bins or sharps
containers

← FULL



RECYCLED BY THE MANUFACTURER
PATENT # 5,887,700 B2



8 890400 0000 8



- When you mix infectious waste and regular solid waste together, you are not permitted to separate them



- Once combined, the entire contents are considered infectious waste!

Contact Information

If you have any questions or comments regarding South Cove's Infectious waste program please contact:

Linda Chu, Human Resources, (617) 521-6701, lchu@scchc.org

Eric Tiberi, COO, (617) 521-6708, etiberi@scchc.org